

ANTIPATHOGENIC EFFICACY OF METHANOLIC LEAF EXTRACT OF CINNAMOMUM TAMALA (BUCH.-HAM.) AND AEGLE MARMELOS (L.) WITH THEIR NUTRITIONAL POTENTIALITY

SUKUMAR DANDAPAT*, MANOJ KUMAR, AMIT KUMAR, M. P. SINHA

Department of Zoology, Ranchi University, Ranchi - 834 008, Jharkhand, INDIA

e-mail: sukumar.msc27@gmail.com

KEYWORDS

Cinnamomum tamala
Aegle marmelos
Antipathogenic
Phytochemical
Nutritional

Received on :
03.03.2013

Accepted on :
15.05.2013

*Corresponding
author

ABSTRACT

Antipathogenic efficacy of methanolic leaf extract of *Cinnamomum tamala* (Buch.-Ham.) and *Aegle marmelos* (L.) through inhibiting the growth of *Staphylococcus aureus* (MTCC 3160), *Salmonella typhi* (MTCC 3216) and *Proteus mirabilis* (MTCC 7837) the causative pathogens of food poisoning, boils, abscesses, wound infection, pneumonia, toxic shock syndrome, typhoid fever, urethritis, cystitis, pyelonephritis and prostatitis has been investigated. All the strains were affected by methanolic leaf extract of *C. tamala* and *A. marmelos* in agar diffusion method and broth dilution method. The MIC values in agar diffusion method were 2.5 mg/mL against *S. aureus* for extract of both plants and 5 mg/mL, 1.25 mg/mL, against *P. mirabilis* for the extract of *C. tamala* and *A. marmelos* respectively. The MIC values in broth dilution method were 2.5 mg/mL against *S. aureus* for both plants extract, 4 mg/mL, 1 mg/mL against *P. mirabilis* and 9 mg/mL, 10 mg/mL against *S. typhi* for *C. tamala* and *A. marmelos* respectively. The nutritional value, phytochemical contents and inorganic substance content of *C. tamala* is higher than *A. marmelos*.

INTRODUCTION

Infectious diseases are disorders caused by pathogenic microorganisms like bacteria, viruses, fungi, protozoa and multicellular parasites. These diseases are also called as communicable or transmissible diseases since they can be transmitted from one person to another via a vector results in the symptoms of disease (Solanki, 2010).

Three common pathogenic bacteria have been tested. *P. mirabilis* is known to cause urethritis, cystitis, pyelonephritis, prostatitis and pneumonia (Todar, 2012). *Staphylococcus* species are predominant among the organisms that are responsible for infective complications following surgical vascular grafts or the implantation of prosthetic devices (De-Lalla, 1999). *Staphylococcus aureus* is a facultative anaerobic, gram positive bacterium, which causes food poisoning and usually grows on the nasal membrane and skin. It is also found in the gastrointestinal and urinary tracts of warm-blooded animals. It also causes boils, abscesses, wound infection, pneumonia, toxic shock syndrome and other diseases (Chesbrough, 2000). Typhoid fever is predominantly caused by *S. typhi* (Crump et al., 2004) and is a global infection (Nagshetty et al., 2010).

Plants are rich in secondary metabolites such as tannins, alkaloids, flavonoids, phenols, etc, which are responsible for therapeutic activities (Rabe and Vnstoden, 2000). The review of literature revealed that considerable contributions have been made on medicinal plants by many workers (Dadsena et al., 2013; Kullu et al., 2013; Kumar et al., 2013; Kumar et

al., 2013a; Mahato et al., 2013; Tabassum et al., 2013; Toppo et al., 2013; Sahu et al., 2013).

Natural products of higher plants may possess a new source of antimicrobial agents with possibly novel mechanisms of action (Barbour et al., 2004; Ahmad and Aqil, 2007). They are effective in the treatment of infectious diseases while simultaneously mitigating many of the side effects that are often associated with synthetic antimicrobials (Iwu et al., 1999). Medicines obtained from plants are relatively safer than synthetic alternative (Iwu et al., 1999; Idu et al., 2007). Therefore, it is of great interest to carry out a screening of these plants in order to validate their use in folk medicine and to reveal the active principle by isolation and characterization of their constituents. Systematic screening of them may result in the discovery of novel active compounds (Tomoko et al., 2002).

Aegle marmelos commonly known as bael, belonging to the family rutaceae and *Cinnamomum tamala* belonging to family lauraceae have been tested against the pathogenic bacteria. These plants are frequently used as folk medicine for various treatments (Chopra et al., 1956; Kirtikar and Basu, 1995; Rao, 2008). The present study is an attempt to evaluate the potentiality of methanolic leaf extract on *Proteus mirabilis*, *Salmonella typhi* and *Staphylococcus aureus*.

MATERIALS AND METHOD

Collection of plant material

The fresh tender leaves of *Aegle marmelos* and *Cinnamomum tamala* were collected from Ranchi (23°21' 0" N LR, 85°20'

0" E L), washed and disinfected with 0.1% HgCl₂ solution and shade dried. Dried material was then powdered in an electric grinder and sieved (Jonani and Sondhi, 2002).

Extract preparation

50g of the powder was subjected to extraction by soxhlet using methanol. The extract obtained was filtered, concentrated in a rotary flash evaporator at 45°C, percentage yield of each extract was calculated and the dried extract was stored in air tight containers at room temperature for further studies.

Phytochemical analyses

Ash content analysis was done following WHO (1998). The amount of crude fiber was determined following Watanables and Olsen (1965). The moisture content was determined in terms of the loss in weight of the plant material on overnight heating at 150°C (Sadasivam and Manickam, 1996). Total phenol was determined by Folin-Ciocalteu reagent, following Ramamoorthy and Bono (2007). The tannins content was quantified as percentage following the procedure and formula given in the quality control methods for medicinal plant materials (WHO, 1998). Aluminium chloride colorimetric method was used with some modifications to determine flavonoids content (Lin and Tang, 2007). Alkaloid was determined by the method used by Helrich (1990). Saponin content was determined following Obdoni and Ochuko (2001).

Nutritive value

Crude fat, carbohydrate and protein were quantified following previously published standard tests (Watanble and Olsen 1965; Jayarama, 2005), and nutritive values were calculated following Nile and Khobragade (2009).

Anti-bacterial analysis

Test Microorganisms

Proteus mirabilis MTCC 7837, *Salmonella typhi* MTCC 3216 and *Staphylococcus aureus* MTCC 3160 used during the present experiment were procured from Hi-media Laboratories (Mumbai, India).

Agar diffusion method

Following Threlfall *et al.* (1999) the agar plates were prepared and wells were made in the plate. Each plate was inoculated with 18 hours old cultures of the selected bacteria and spread evenly on the plate. After 20 minutes, the wells were filled with different concentrations of samples. The control wells were filled with Gentamycin along with solvent. All the plates were incubated at 37°C for 24h and the diameter of inhibition zones were noted.

Broth dilution method

As proposed by Walker (2000) the tubes containing the culture media were prepared, autoclaved and respective concentrations of the samples were added. Each tube was inoculated with 18 hours old cultures (100iL, 10⁴cfu). A control tube with inoculums and without any sample was prepared along with a sterile media tube as blank. All the tubes were incubated at 37°C on a shaker with 140 rpm for 24h; the growth and hence the MIC was measured at 660nm.

RESULTS AND DISCUSSION

Phytochemical profile

The results on phytochemical analyses of *C. tamala* and *A. marmelos* leaf have been represented in Fig. 1, 2 and 3. The result reveal that moisture content is higher in *A. marmelos* leaf than *C. tamala* and ash content is lower in *A. marmelos* leaves than *C. tamala* leaves.

Physicochemical analysis from leaves of various medicinal plants has been reported by various workers from time to time. Indrayan *et al.* (2005) reported 8.20 g/100g ash, 57.90 g/100g moisture and 7.20g/100g crude fiber in *A. hetrophyllus* leaves. Nasiruddin *et al.* (2012) detected 2.84 ± 0.04%, 1.33 ± 0.02%, 3.50 ± 0.03% total ash, 91.60 ± 0.20%, 82.90 ± 0.74%, 85.05 ± 0.50% moisture and 0.94 ± 0.06%, 3.11 ± 0.05%, 2.41 ± 0.05% crude fiber in *Rumex crispus*, *Medicago denticulate* and *Taraxicum officinale* respectively.

The amount and composition of ash remaining after combustion of plant material varies considerably according to the part of the plant, age, treatment etc. Ash usually represents the inorganic part of the plant (Vermani *et al.*, 2010). Nutritionally, fiber is beneficial to human body, since it has been reported that food fiber aids absorption of trace elements in the gut (Kelsay, 1981) and reduce absorption of cholesterol (Le - Veille and Sanberlich, 1966). Fiber aids bowel movement of gut (Abolaji *et al.*, 2007). Aravind *et al.* (2013) reported, fiber of *Carica papaya* is able to bind cancer-causing toxins in the colon and keep them away from the healthy colon cells. The fibers provide synergistic protection for colon cells from free radical damage to their DNA. *C. tamala* and *A. marmelos* leaves contain higher amount of ash and crude fiber (Fig. 1) compared to the above plants thus *C. tamala* and *A. marmelos* leaf are likely to contain higher amount of inorganic constituent and dietary fibers.

The results on phytochemical analysis of the leaf samples of *C. tamala* and *A. marmelos* is presented in Fig - 3. The result revealed that polyphenols is highest (16.7 ± 0.7 g/100g), flavonoid is lowest (1.0 ± 1.01 g/100g) in *C. tamala* and polyphenols is highest (6.7 ± 0.61 g/100g), alkaloid is lowest (2.3 ± 0.42 g/100g) among all the studied phytochemicals. Aliyu *et al.* (2008) reported 0.110 ± 0.002 g/100g, 0.966 ± 0.030 g/100g, 1.440 ± 0.002 g/100g, 7.270 ± 0.009 g/100g and 2.600 ± 0.200 g/100g alkaloids, 8.000 ± 0.280g/100g, 8.766 ± 0.020g/100g, 16.30 ± 0.042g/100g, 18.23 ± 0.040 g/100g and 9.466 ± 0.060 g/100g flavonoids, 0.533 ± 0.020 g/100g, 2.500 ± 0.014 g/100g, 0.900 ± 0.020 g/100g, 2.320 ± 0.001 g/100g and 1.066 ± 0.020 g/100g saponins, 0.566




Figure 1: Physicochemical composition of *C. tamala* and *A. marmelos* leaf in g/100g (M ± SD; n = 3).

± 0.010 g/100g, 1.250 ± 0.009 g/100g, 0.520 ± 0.200 g/100g, 1.030 ± 0.014 g/100g and 1.140 ± 0.001 g/100g phenols in *Anchomanes difformis*, *Anisopus mannii*, *Pavetta crassipes*, *Stachytarpheta angustifolia* and *Vernonia blumeoides* respectively. Manikandan et al. (2010) reported 10.0 mg/g and 13.0 mg/g tannin in *Ruellia tuberosa* L. and *Dipteracanthus patulus* (Jacq.) respectively. Soladoye and Chukwuma (2012) reported tannin (4.98%) in *Cissus populnea*. Khan et al. (2011) reported tannin content 15.75% in *M. rubicaulis*, 14.16%, *W. fruticosa*, 13.4% in *C. grata*, 12.33% in *V. cotinifolium*, 11.2% in *E. hirta*, 10.56% in *B. Papyrifera* and 10.2% in *P. harmala*.

The total phenolic content of *Cinnamomum tamala* and *Aegle marmelos* 16.7 ± 0.7 g/100g and 6.7 ± 0.42 g/100g respectively have been found highest among most of the plants studied. Tannins, alkaloids, saponins, flavonoids, and sterols have been found active against several pathogenic bacteria (Kennedy and Wightman, 2011, Choudhury et al., 2013). Tannins form irreversible complexes with proline rich protein resulting in the inhibition of cell wall synthesis (Mamtha et al., 2004).

Flavonoids inhibit several enzymes, chelate certain metal cations, affect protein phosphorylation (Middleton and Kandaswami, 1994) and have variety of effects on membrane-linked processes (Smith, 1996) including the enhancement of metal-induced lipid peroxidation (Sakihama et al., 2002). Alkaloids possess anti-oxidizing effects, thus reduces the nitrate generation which is useful for protein synthesis, suppresses the transfer of sucrose from stomach to small intestine. Isaac and Chinwe (2001) reported that alkaloids are responsible for the antibacterial activity.

Nutrition potentiality

The result of nutritional potentiality of *C. tamala* and *A. marmelos* leaves have been represented in fig – 2 and table - 1. The results reveal that carbohydrate content is higher in *A. marmelos* leaves (10.5 ± 0.3 g/100g) than *C. tamala* leaf (9.5 ± 0.5 g/100g) and fat content is lower in *A. marmelos* leaves (1.7 ± 0.5 g/100g) than *C. tamala* leaves (6.0 ± 0.5 g/100g).

Indrayan et al. (2005) reported 19.70% carbohydrate, 5.70% protein and 2.50% crude fat in *A. hetrophyllus* leaves. Bukhsh et al. (2007) reported $18.9 \pm 4.2\%$, $16.9 \pm 1.1\%$, $15.9 \pm 1.3\%$ carbohydrate and $21.87 \pm 4.7\%$ crude protein in *Carthamus oxyacantha*, *Eruca sativa* and *Plantago ovate* leaves respectively. The fat content was $6.6 \pm 1.3\%$ in *Eruca*




Figure 2: Composition of fat, protein and carbohydrate and nutritive value from *C. tamala* and *A. marmelos* leaf in g/100g ($M \pm SD$; $n = 3$).

sativa leaves but fat was not found in *Carthamus oxyacantha* and *Plantago ovate* leaves (Bukhsh et al., 2007). Nasiruddin et al. (2012) reported $1.82 \pm 0.03\%$, $5.99 \pm 0.02\%$, $2.74 \pm 0.01\%$ crude protein and $0.30 \pm 0.01\%$, $0.14 \pm 0.03\%$, $0.21 \pm 0.02\%$ crude fat in *Rumex crispus*, *Medicago denticulate* and *Taraxicum officinale* respectively.

Since carbohydrate constitutes a major class of naturally occurring organic compounds that are essential for the maintenance of animal life (Ebun-Oluwa and Alade, 2007). Proteins contain amino acids utilized by the cells of the body to synthesize all the numerous proteins required for the function of the cell and also to furnish energy (Robinson, 1978). Due to low level of crude fat in the leaves of *A. marmelos* and *C. tamala*, the leaves can be consumed in diet of those people suffering from overweight or obesity (Nasiruddin et al., 2012).

The calculated nutritional value is higher (143.5 ± 0.53 Kcal/100g) in *C. tamala* than *A. marmelos* (82.5 ± 0.74 Kcal/100g). Nasiruddin et al. (2012) reported total energy 21.15 Kcal, 55.05 Kcal and 48.46 Kcal in *Rumex crispus*, *Medicago denticulate* and *Taraxicum officinale* respectively. Indrayan et al. (2005) reported 124.10 cal/100 g nutritive values of *A. hetrophyllus* leaves. The nutritional values of indigenous fruits and vegetables such as *Cucumis sativus*, *Pangium edule*, *Brassica oleraceae*, *Spinacia oleraceae*, *Sinapis alba* have been reported as 15 kcal, 227 kcal, 22 kcal, 29 kcal, 34 kcal respectively (Hoe and Siang, 1999).

Table 1: Nutritional value of *C. tamala* and *A. marmelos* ($M \pm SD$; $n = 3$).

Nutritional value	<i>C. tamala</i>	<i>A. marmelos</i>
	143.5 ± 0.53	82.5 ± 0.74
	Kcal/100g	Kcal/100g

Since *C. tamala* and *A. marmelos* leaves contain high amount of carbohydrate, protein, fat (Fig. 2) and nutritional value comparing with the above plants, thus leaves of *C. tamala* and *A. marmelos* can be used as fodder.

Antibacterial analysis

The pathogenic efficacy of methanolic extract of *C. tamala* and *A. marmelos* leaves were quantitatively assessed on the basis of zone of inhibition (ZOI) in mm (Table 2) following the agar disk diffusion method and minimum inhibitory concentration by broth dilution method. The test organisms were inoculated with standard antibiotic: gentamycin to compare the efficacy of leaf extract for their microbial properties (Table 3). In the present investigation the extracts were found




Figure 3: Phytochemicals from *C. tamala* and *A. marmelos* leaf in g/100g ($M \pm SD$; $n = 3$).




Figure 4: (a, b, c) ZOI for the methanolic extract of *A. marmelos*; Fig 4:(d, e,f) ZOI for the methanolic extract of *C. tamala*; (g,h, i) ZOI for the gentamycin.

to be effective against all the pathogens. The ZOI observed for the methanolic extract and gentamycin using agar diffusion method is represented in Fig. 4a, 4b, 4c, 4d, 4e, 4f and Fig. 4g, 4h, 4i respectively. The broth dilution method showed more pronounced antimicrobial activity through 100% inhibition of all the pathogens in the range of 1.25-10mg/mL concentration (Fig. 5a, 5b, 5c, 5d, 5e and 5f). The minimum inhibitory concentration (MIC) obtained by broth dilution method for *P. mirabilis*, *S. aureus* and *S. typhi* were in the range of 1mg/mL-10 mg/mL. Kothari *et al.* (2011) worked out on different extract of *A. marmelos* and found $10 \pm 0.3 \text{ mm} - 22 \pm 0.6 \text{ mm}$ zone of inhibition against *S. aureus*, *S. typhi*, *P. mirabilis* and other pathogenic bacteria species respectively and also said that methanolic extract was more effective than other extracts. Essawi and Srouns (2000) reported that methanolic leaf extract was more effective compared to chloroform and aqueous extract because of chemical constituents which are either polar or non polar and can be effectively extracted only through the organic solvent medium. *Cinnamomum tamala* possess antibacterial activity due to the presence of certain phenolic compound such as cinnamic aldehyde and such as eugenol and cinnamic acid (Baratta *et al.*, 1998). An important characteristic of leaf extract and their

Table 2: The zone of inhibition and MIC (in mm) of methanolic leaf extract of *C. tamala* and *A. marmelos*.

Concentration (mg/mL)	<i>C. tamala</i>			<i>A. marmelos</i>		
	<i>P. mirabilis</i>	<i>S. aureus</i>	<i>S. typhi</i>	<i>P. mirabilis</i>	<i>S. aureus</i>	<i>S. typhi</i>
0.18	-	-	-	-	-	-
0.36	-	-	-	-	-	-
0.612	-	-	-	-	-	-
1.25	-	-	-	6	-	-
2.5	-	6	-	9	4	-
5	10	6	-	10	5	-
MIC(mg/mL)	5	2.5	-	1.25	2.5	-

Table 3: The zone of inhibition and MIC (in mm) of Gentamycin against the test organism.

Concentration (mg/mL)	<i>S. typhi</i>	<i>S. aureus</i>	<i>P. mirabilis</i>
0.025	2	13	9
0.05	13	18	13
0.10	16	21	18
0.20	21	25	21
0.40	25	27	25
0.80	27	34	27
MIC(mg/mL)	00.25	00.25	00.25

components is their hydrophobicity, which enable them to partition the lipids of the bacterial cell membrane and mitochondria, disturbing the cell structure and rendering them




Figure 5: (a) and (b) Inhibition % of *S. aureus*, (c) and (d) Inhibition % of *S. typhi*, (e) and (f) Inhibition % of *P. mirabilis* in broth dilution method for Methanolic leaf extract of *C. tamala* and *A. marmelos* respectively.

more permeable (Sikkema *et al.*, 1994). Extensive leakages from bacterial cells or exits of critical molecules and ions will lead to death (Denyer and Hugo, 1991). The antibacterial activity *A. marmelos* leaf extract is due to presence of active phenolic compound eugenol and cuminaldehyde because these compounds have already shown their activity against various bacterial strains (Duke, 1992) the mechanism of action may be the blockage of protein synthesis either at transcription or at translation level and inhibition of peptidoglycan synthesis at membrane level (Rajan and Jeevagangai, 2009).

The present study suggests antibacterial property of *Cinnamomum tamala* and *A. Marmelos* leaf extract, which inhibits the growth of pathogenic bacteria *S. aureus*, *S. typhi* and *P. mirabilis* causative agent food poisoning, boils, abscesses, wound infection, pneumonia, toxic shock syndrome, typhoid fever and urethritis, cystitis, pyelonephritis, prostatitis and pneumonia disease, and can be used as new drug for therapy.

ACKNOWLEDGEMENT

The authors acknowledge the facilities provided by the Department of Zoology, Ranchi University, Ranchi.

REFERENCES

Abolaji, O. A., Adebayo, A. H and Odesanmi, O. S. 2007. Nutritional

Qualities of Three Medicinal Plant Parts (*Xylopiya aethiopicpa*, *Blighia sapida* and *Parinari polyandra*) commonly used by Pregnant Women in the Western Part of Nigeria. *Pak. J. Nut.* **6(6)**: 665-668.

Ahmad, I. and Aqil, F. 2007. In vitro efficacy of bioactive extracts of 15 medicinal plants against ESBL-producing multidrug-resistant enteric bacteria. *Microbiol. Res.* **162**: 264-275.

Aliyu, A. B., Musa, A. M., Oshanimi, J. A., Ibrahim, H. A. and Oyewale, A. O. 2008. Phytochemical analysis and mineral elements composition of some medicinal plants of Northern Nigeria. *Niger. J. Pharmaceutical Sci.* **7(1)**: 119-125.

Aravind, G., Bhowmik, D., Duravel, S. and Harish, G. 2013. Traditional and Medicinal Uses of *Carica papaya*. *J. Med. Plnt. Stud.* **1(1)**: 7-15.

Baratta, M. T., Dorman, H. J., Deans, S. G., Figueiredo, A. C., Barroso, J. G. and Ruberto, G. 1998. Antimicrobial and antioxidant properties of some commercial essential oils. *Flav. Fragr. J.* **13**: 235-244.

Barbour, E. K. A., Sharif, M., Sagherian, V. K., Habre, A. N., Talhouk, R. S. and Talhouk, S. N. 2004. Screening of selected indigenous plants of Lebanon for antimicrobial activity. *J. Ethnopharmacol.* **93**: 1-7.

Bukhsh, E., Mallik, S. A. and Ahmaddh, S. S. 2007. Estimation of Nutritional value and trace element content of *Carthamus oxyacantha*, *Eruca sativa* and *Plantago ovata*. *Pak. J. Bot.* **39(4)**: 1181-1187.

Cheesbrough, M. 2000. Medical Laboratory Manual for Tropical Countries. Microbiology, Linacre house, Jordan Hill Oxford. p. 260.

Chopra, R. N., Nayar, S. L. and Chopra, I. C. 1956. Glossary of Indian Medicinal Plant, fifth Edition, IBH publication (P) Ltd, New Delhi. pp. 8-14.

Choudhury, S., Sharan, L. and Sinha, M. P. 2013. Phytochemical and antimicrobial standardization of the methanolic leaf extracts of *Murraya Koenigii* Linn. *Aech. Des. Sci.* **66(3)**: 67-80.

Crump, J. A., Okoth, G. O., Slutsker, L., Ogaja, D. O., Keswick, B. H. and Luby, S. P. 2004. Effect of point-of-use disinfection, flocculation and combined flocculation/disinfection on drinking water quality in western Kenya. *J. Appl. Microbiol.* **97**: 225-231.

Dadsena, R., Sahu, N. K., Agrawal, S. and Kumar, A. 2013. Phytochemical analysis of three endangered plants (*Costus speciosus*, *Gloriosa superba* Linn and *Rauvolfia serpentina* (Linn) Benth from Kanker district of Chhattisgarh, India. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 655-659.

De-Lalla, F. 1999. Antimicrobial chemotherapy in the control of surgical infectious complications. *J. Chemotherap.* **11**: 440-445.

Denyer, S. P. and Hugo, W.B. 1991. Biocide-included damage to the bacterial cell membrane. The society for applied bacteriology, Technical series No.-27. Oxford Blackwell scientific publication, Oxford. pp. 171-188.

Duke, J. A. 1992. Hand book of biologically active phytochemicals and their activities. 48ED. CRC press. Boca Raton, Florida. p. 111.

Ebun-Oluwa, P.O and Alade, A.S. 2007. Nutritional potential of Berlandier Nettle spurge (*Jatropha cathatica*) seed. *Pak. J. Nutr.* **6**: 345-348.

Essawi, T. and Srour, M. 2000. Screening some Palestinian medicinal plants for antibacterial activity. *J. Ethnopharmacol.* **70**: 343-349.

flavonoids on mammalian biology: implications for immunity, inflammation and cancer In: Harborne, J.B.(eds). The Flavonoids Advances in Research Science. Chapman and Hall, London, pp. 619-652.

Helrich, K. 1990. Official Methods of Analysis of the Association of Official Analytical Chemists. AOAC, Inc. (15th ed). Virginia, USA, pp. 425-497.

Hoe, V. B. and Siong, K. H. 1999. The nutritional value of indigenous

- fruits and vegetables in Sarawak. *Asi. Paci. J. Clin. Nutr.* **8(1)**: 24-31.
- Idu, M., Omogbai, E. K. I., Aghimien G. E. I., Amaechina, F., Timothy, O. and Omonigho, S. E. 2007.** Preliminary phytochemistry, antimicrobial properties and acute toxicity of *Stachytarpheta jamaicensis* (L.) Vahl. *Leaves. Tre. Med. Res.* **2**: 193-198.
- Indrayan, A. K., Sharma, S., Durgapal, D., Kumar, N. and Kumar, M. 2005.** Determination of nutritive value and analysis of mineral elements for some medicinally valued plants from Uttaranchal. *Crr. Sci.* **89(7)**: 1252-1255.
- Isaac, O. O and Chinwe, J. A. 2001.** The phytochemical analysis and antibacterial screening of extract of *Tetrecarpidium conophorum*. *J. Chem. Soc. Nig.* **26(1)**: 53-55.
- Iwu, M. W., Duncan, A. R. and Okunji, C. O. 1999.** New antimicrobials of plant origin In: Perspectives on new Crops and new Uses, eds. J. Janick, ASHS Press, Alexandria, VA. Pp. 457-462.
- Jayaram, J. 2005.** Laboratory manual in Biochemistry, New age international (P) Ltd. **24**: pp 75 – 78.
- Jonani, G. K. and Sondhi, S. M. 2002.** Determination of minerals elements in some Ayurvedic bhasmas used for the cure of various elements. *Phytother. Res.* **16**: 774-777.
- Kelsay, J. L. 1981.** Effects of diet fiber on bowel function and trace mineral balances of human subjects. *Cereal Chem.* **58**: 2-5.
- Kennedy, D. O. and Wightman, E. L. 2011.** Herbal extracts and phytochemicals: Plant secondary metabolites and enhancement of human brain function. *Adv. Nutrit.* **2**: 32-50.
- Khan, A. M., Qureshi, R. A., Ullah, F., Gilani, S. A., Nosheen, A., Sahreen, S., Laghari, M. K., Laghari, M. Y., Rehman, S. U., Hussain, I. and Murad, W. 2011.** Phytochemical analysis of selected medicinal plants of Margalla Hills and surroundings. *J. Med. Plant Res.* **5(25)**: 6055-6060.
- Kirtikar, K. R. and Basu, B. D. 1995.** Indian medical plant, second Edition, International book publication, Mumbai, India. pp. 499-502.
- Kothari, S., Mishra, V., Bharat, S. and Tonpay, S. D. 2011.** Antimicrobial activity and phytochemical screening of serial extracts from leaves of *Aegle marmelos* (Linn.). *Acta Pol. Pharma. Drug Res.* **68(5)**: pp. 687-692.
- Kullu, A. R., Tabassum, W. and Sinha, M. P. 2013.** Effect of *Psidium guajava* aqueous extracts on haematological profile and serum lipid variables of albino rat. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 743-746.
- Kumar, A., Kumar, M., Dandapat, S. and Sinha, M.P. 2013.** Antioxidant activity and pharmacological screening of *Tinospora cordifolia*. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 689-693.
- Kumar, M., Kumar, A., Dandapat, s. and Sinha, M. P. 2013.** Phytochemical screening and antioxidant potency of *Adhatoda vasica* and *Vitex negundo*. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 723-730.
- Le Veille, G. and Sanberlich, H.E. 1966.** Mechanism of the cholesterol-dressing effect of pectin in the cholesterol fed rat. *J. Nutr.* 209-214.
- Lin, J. Y and Tang, C. Y. 2007.** Determination of total phenolic and flavonoids contents in selected fruits and vegetables, as well as their stimulatory effects on mouse spleenocyte proliferation. *Food Chem.* **101(1)**: 140-147.
- Mahato, S., Mehta, A. and Roy, S. 2013.** Studies on antibacterial effects of bark, seed and callus extracts of *Holarrhena antidysenterica* Wall. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 717-721.
- Mamtha, B., Kavitha, K., Srinivasan, K. K. and Shivananda, P. G. 2004.** An in vitro study of the effect of *Centella asiatica* (Indian pennywort) on enteric pathogens. *Ind. J. Pharmacol.* **36(1)**: 41.
- Manikandan, A., Victor, D. and Doss, A. 2010.** Evaluation of biochemical contents, nutritional value, trace elements, SDS-PAGE and HPTLC profiling in the leaves of *Ruellia tuberosa* L. and *Dipteracanthus patulus*(Jacq.). *J. Chem. Pharm. Res.* **2(3)**: 295-303.
- Middleton, E. and Kandaswami, C. 1994.** The impact of plant flavonoids on mammalian biology: Implications for immunity, inflammation and cancer in harbore, J.B. (eds.). *The flavanoids Advances in Research Sciences.* Chapman and Hall, London. pp. 619-652.
- Nagshetty, K., Channappa, S.T. and Gaddad, S. M. 2010.** Antimicrobial susceptibility of *Salmonella typhi* in India. *J. Infect. Dev. Count.* **4(2)**: 70-73.
- Nasiruddin., Masood, T., Arif, M., Shah, S. S., Azhar, N. and Saifullah. 2012.** Nutritional content of some medicinal herbs of Peshawar district, Pakistan. *Sarhad. J. Agri.* **28(4)**: 335-339.
- Nile, S. H. and Khobragade, C. N. N. 2009.** Determination of nutritive value and mineral elements of some important medicinal plants from western part of India. *J. Med. plants.* **8(5)**: 79-88.
- Obadona, B. O., Ocheke, P. O. 2001.** Phytochemical studies and comparative efficacy of the crude extract of some homeostatic plants in Edo and Delta states of Nigeria. *Glob. J. Pure and Appl. Sci.* **86**: 2003-2008.
- Rabe, T. and Vanstoden, J. 2000.** Isolation of an antimicrobial sesquiterpenoid from *Warbugie salutaris*. *J. Pharmacol.* **93**: 171-174.
- Rajan, S. and Jeevagangai, T. J. 2009.** Studies on the antibacterial activity of *Aeglemarmelos*–fruit pulp and its preliminary phytochemistry. *J. Bas. Appl. Biol.* **3(1&2)**: 76-81.
- Ramamoorthy, P. K and Bono, A. 2007.** Antioxidant activity, total phenolic and flavonoids content of *Morinda citrifolia* fruit extracts from various extraction processes. *J. Engg. Sci. Tech.* **2(1)**:70-80.
- Rao, G. V. 2008.** The chemical constituents and biological studies of *Chloroxylon swietenia*. *Ind Drug.* **45(1)**: 5-15.
- Robinson, H. 1978.** Fundamentals of normal nutrition, 3rd Edition, Macmillon Publishing Co., Inc. ISBN 02-979590. pp. 41-125, 271-284.
- Sadasivam, S and Manickam, A. 1996.** Biochemical methods. 2nd (ed). New Age. International Limited Publishers, New Delhi, India. pp. 159- 160.
- Sahu, P. R. and Sinha, M. P. 2013.** Screening of antibacterial activity of crude leaf extracts of *Cassia tora* on UTI pathogens. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 735-738.
- Sakihama, Y. Cohen, M. F. Grace, S. C and Yamasaki, H. 2002.** Plant phenolic antioxidant and prooxidant activities: Phenolics-induced oxidative damage mediated by metals in plants. *Toxicol.* **177**: 67-80.
- Sikkema, J., Bont de, J. A. M and Poolman, B. 1994.** Interactions of cyclic hydrocarbons with biological membranes. *J. Biol. Chem.* **269**: 1022-8028.
- Smith, C. J. 1996.** Accumulation of phytoalexins: Defence mechanism and stimulus response system. *New Phytol.* **132(1)**: 1-45.
- Soladoye, M. O. and Chukwuma, E. C. 2012.** Quantitative phytochemical profile of the leaves of *Cissus populnea* Guill. and Perr. (Vitaceae) –an important medicinal plant in central Nigeria. *Arch. App. Sci. Res.* **4(1)**: 200-206.
- Solanki, R. 2010.** Some medicinal plants with antibacterial activity. *Int. J. Comprehens. Pharm.* **4(10)**: 1-4.
- Tabassum, W., Kullu, A. R. and Sinha, M. P. 2013.** Effects of leaf extracts of *Moringa oleifera* on regulation of hypothyroidism and lipid profile. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 665-669.
- Threlfall, E. J., Fisher, I. S. T., Ward, L., Tschape, H. and Gernersmidt, P. 1999.** Harmonization of antibacterial susceptibility testing for *Salmonella*: Result of a study by 18 national reference laboratories

within the European Union-funded Enter-Net group. *Microbiol. Drug Resist.* **5**: 195-199.

Todar, K. 2012. Online textbook of Bacteriology. <http://www.textbookofbacteriology.net>

Tomoko, N., Takashi, A., Hiromu, T., Yuka, I., Hiroko, M., Munekazu, I., Totshiyuki, T., Tetsuro, I., Fujio, A., Iriya, I., Tsutomu, N. and Kazuhito, W. 2002. Antibacterial activity of extracts prepared from tropical and subtropical plants on methicillin-resistant *Staphylococcus aureus*. *J. Health Sci.* **48**: 273-276.

Toppo, K. I., Gupta, S., Karkun, D., Agrawal, S. and Kumar, A. 2013. Antimicrobial activity of *Sphagneticola trilobata* (L.) Pruski, against some human pathogenic bacteria and fungi. *The Bioscan.* **8(2): Supplement on Medicinal Plants.** 695-700.

ukhsh, E., Malik, S. A. and Ahmadh, S. S. 2007. Estimation of nutritional

value and trace element content of *Carthamus oxyacantha*, *Eruca sativa* and *Plantago ovata*. *Pak. J. Bot.* **39(4)**: 1181-1187.

Vermani, A., Prabhat, N. and Chauhan, A. 2010. Physico-Chemical Analysis of Ash of Some Medicinal Plants Growing in Uttarakhand, India. *Nature. Science.* **8(6)**: 88-91.

Walker, R. D. 2000. Antimicrobial susceptibility testing and interpretation of results. *In: Prescott, I.F., Baggot, I.D., Walker, R. D., Ames, I. A. (3rd Ed).* Antimicrobial Therapy in Veterinary Medicine. Iowa State University Press. pp. 12-26.

Watanble, F. S. and Olsen, S. R. 1965. Test for ascorbic acid method for determining phosphorus in water and sodium bicarbonate extract of soil. *Proc. Soil. Sci. Soc. Am.* **29**: 677 – 678.

WHO 1998. Quality control methods for medicinal plant materials. Library Cataloguing in Publication data. p. 44.

