

ROLE OF SECONDARY METABOLITES ON BIOCONTROL POTENTIALITIES OF NATIVE RHIZOBACTERIAL ISOLATES AGAINST RHIZOCTONIA SOLANI

ADAM KAMEI,* S. DUTTA AND S. NANDI

Department of Plant Pathology, AICRP on Vegetable Crops, Directorate of Research, BCKV, Kalyani - 741235, West Bengal
e-mail: kameiadam.03@gmail.com

KEYWORDS

Rhizobacteria
Rhizoctonia solani
Rice
Secondary metabolites
predictors

Received on :

04.10.2013

Accepted on :

29.02.2014

*Corresponding
author

ABSTRACT

Sixty two isolates of rhizobacteria were isolated from rhizosphere of rice crop, chilli, cabbage and potato collected from Gangetic Alluvial regions of West Bengal. Rhizobacterial isolates were evaluated *in-vitro* by dual culture method. Ten of which were found to be potent bioagents. The isolates were identified as *Pseudomonas aeruginosa* based on biochemical and molecular identification techniques. Studies revealed that the mycelial growth of *R. solani* was inhibited up to 1.9 (cm) by PTR-3 and were found to exhibit antagonism of over 68.9% which is followed by PCF-3(65.6%). Isolate PTR-3 having higher level of chitinolytic (17.2 pmol/s) and proteolytic (halo zone 2.9 cm) activity was highly effective against *R. solani*. Among the ten antagonistic rhizobacteria, highest level of salicylic acid(0.54), siderophore(3.9 μ mole benzoic acid /ml) and HCN production was noticed by PTR-1 isolate. Whereas, highest level of IAA(0.01039 μ g/ml) and Phosphate solubilisation(17.3 μ g/ml) activity were noticed by PTR-4 and PTR-3 isolates respectively. Extracellular enzymes activity of protease and chitinase were found to be highly and positively correlated with the ability to antagonize *R. solani*. Step wise regression analysis revealed that the combination of biochemical variables chitinase and protease activity jointly contributing 98% of variability of antagonistic potentiality of rhizobacteria against, *R. solani* causing sheath blight disease of rice. These variables may be used as predictors of biocontrol potentiality of rhizobacteria against destructive soil-borne fungal pathogen *Rhizoctonia solani*.

INTRODUCTION

The beneficial effects of plant growth promoting rhizobacteria (PGPR) have been exploited in many areas including biofertilizers, microbial rhizo-remediation and biopesticides (Adesemoye *et al.*, 2008). PGPR group offers an effective means of antagonism against sheath blight pathogen, *R. solani* (Luo Jin Yan *et al.*, 2005). Secondary metabolites produced by fluorescent Pseudomonads have been reported antifungal activity inhibiting *R. solani* (Mina *et al.*, 2013). Rice sheath blight pathogen, *R. solani* suppressed by non-fluorescent PGPR like *Bacillus* sp was previously attributed to the production of chitinase (Krishnaveni, 1991) and other antifungal metabolites. A large number of bacterial strains were found to possess the ability to protect rice plants from sheath blight disease (Vasantha Devi *et al.*, 1989) and these were identified through dual plate assays. In recent years, fluorescent Pseudomonads have drawn worldwide attention as they produce secondary metabolites such as siderophore, antibiotics, volatile compounds HCN, enzymes and phytohormones (Reddy *et al.*, 2007). PGPR have been considered as an alternative to agrochemicals for controlling plant diseases as plant pathogens developed resistance against pesticides. Therefore, in the present investigation attempts were made to studies the secondary metabolites as bio-control potentiality produced by fluorescent pseudomnads against *Rhizoctonia solani*

MATERIALS AND METHODS

Isolation of Rhizobacteria

The soil samples were collected from rhizosphere of healthy plants of different rice field grown crops. The bacterial suspension thus obtained was used for the isolation of rhizobacteria by dilution plating method King's B medium (Johnson and Curl, 1977). After 48 hours of growth at 28°C (\pm 2°C), morphologically different colonies were picked up and purified.

Maintenance of rhizobacterial isolates and fungal cultures Rhizobacteria isolates

For short period maintenance, the individual bacterial colony was picked up and streaked on yeast glucose dextrose agar, incubate at 28°C (\pm 2°C) and kept at 4°C. Whereas in long term storage single colony was inoculated into 10 ml of nutrient broth, incubate at 28°C for 24 hours followed by shaking at 200 rpm, the broth was mixed with 50% glycerol solution at 3:1 and kept in cryobox at -80°C.

Pathogen fungal cultures

The fungal cultures *Rhizoctonia solani* obtained from Department of Plant Pathology, BCKV was maintained on Potato Dextrose Agar and preserved in the refrigerator at 4°C. Fungal cultures were examined in order to test their pathogenicity at regular intervals weekly.

In vitro antagonistic activity against soil borne fungal pathogens

The antagonistic effects of the rhizobacterial isolates were observed against *R. solani* by dual culture technique (Vincent, 1947). For each treatment three replications were used. Percent inhibition over control was calculated by using the formula.

$$I = \frac{C - T}{C} \times 100$$

I = Per cent inhibition of mycelium

C = Growth of mycelium in control

T = Growth of mycelium in treatment

Characterization of Rhizobacteria

The antagonistic rhizobacterial isolates were characterized on the basis of their morphological (cell shape, cell arrangement), cultural (colony type, pigment production) and biochemical (arginine dihydrolase activity, H₂S production, nitrate reduction etc.) identification scheme of Stolp and Gadhari, 1981 and for *Pseudomonas* sp. (Bossis, 1995), biovars (Bossis *et al.*, 2000) and seedling vigour method as described (ISTA, 1966).

Qualitative assay of HCN, IAA, SA production, phosphate solubilization and siderophore production by the bacterial isolates

HCN [Lorck (1948) modified by Alstrom (1989)], IAA (Gordon and weber, 1951), Salicylic acid (Meyer *et al.*, 1992), Siderophore production (Reeves, 1983) and Phosphate solubilization (by Ammonium phosphor-molybdate method Jeon *et al.*, 2003) was done by spectrophotometric quantitative assay following the methods as mentioned for each assay.

Quantitative and Qualitative assay of different enzyme activities

Protease (Smibert and Krieg, 1994), amylase, cellulase, pectinase and chitinase (Vessey and Pegg, 1973) activity of different rhizobacterial isolates was observed accordingly.

RESULTS AND DISCUSSION

Isolation of rhizospheric bacteria of rice, chilli, cabbage and potato were made isolated on King's B medium, showed 62 rhizospheric isolates present on these crops. Twenty five rhizobacterial isolates were obtained from rice, 12 from chilli, 9 from cabbage and 15 from potato. All the bacteria were screened for their *in vitro* antagonistic activity against soil borne *Rhizoctonia solani*. Among these, ten isolates showed potential antagonistic activity against all the tested soil-borne fungal *R. solani*.

In-vitro antagonism of rhizospheric bacteria against fungal plant pathogen and biochemical identification

Perusal of the data in Table 1 and 2 indicated that all the ten rhizobacterial isolates showed antagonistic effect against *R. solani* under *in vitro* condition and inhibited the vegetative growth of the fungus at varied level. Among the isolates, PTR-3 was found to be the most effective isolate giving an inhibition zone of 1.9 cm and also caused 68.9 % growth inhibition against *R. solani* followed by PCF-3 (1.7 cm) and PTR-1 (1.5 cm). The antagonistic nature of rhizospheric bacteria against *R. solani* was also reported by Adhikari *et al.* (2013). Tripathi

and Johri (2002) observed *in vitro* inhibition of *Colletotrichum dermatium*, *R. solani* and *Sclerotium rolfsii* by fluorescent pseudomonads. Based on the biochemical characterization following the identification scheme of Bossis *et al.* (2000) revealed that all the isolates were *Pseudomonas aeruginosa* as depicted in Table 3.

Studies on protease, amylase, cellulase, pectinase and chitinase activity of different rhizobacterial isolates

Production of chitinolytic enzyme was also analyzed because this is an important means of fungal inhibition. Chitinase was detected in all the ten isolates and isolate PTR (17.2) p mol /s) expressed maximum activity of chitinase followed by PCF-3 (15.6 p mol /s). Rhizobacterial isolates could produce halo zones on skim milk agar and PPF-7 from potato rhizosphere exhibits higher protease activity in terms of production of halo zones (3 cm) on skim milk agar followed by PTR-3 (2.9 cm) (Table 4). But cellulase, amylase, pectinase assay were found to be negative in all the tested rhizospheric isolates.

Role of secondary metabolites and enzyme activities on antagonistic potentiality of rhizobacteria against *R. solani*

Among the rhizobacterial isolates though PTR-1 produced highest HCN (0.0834), salicylic acid (0.54 mg/ml), siderophore (3.9 mM benzoic acid/ml) but PTR-4 produced maximum IAA

Table 1: *In vitro* antibiosis as inhibition zone in cm produced by the effective bacterial isolates against soil borne plant pathogens

Antagonistic Bacterial Isolate	Soil Rhizosphere	Pathogenic fungi (<i>Rhizoctonia solani</i>) Inhibition zone (in cm)
PTR-1	Rice	1.5 ^c
PCF-3	chilli	1.7 ^b
PTR-3	Rice	1.9 ^a
PPF-2	Potato	1.1 ^f
PPF-7	Potato	1.2 ^e
PFC-1	cabbage	1.4 ^d
PTR-4	Rice	1.4 ^d
PCF-7	Chilli	0.9 ^g
PFC-2	Cabbage	0.7 ^h
PFC-7	Cabbage	0.9 ^g

Data bearing same letter are not significantly different on the basis of DMRT at 5% level of significance

Table 2: Growth rate of mycelium of fungal pathogen against rhizobacterial isolate after 72 hrs

Sl. No.	Isolates	Pathogen	Growth rate of mycelium of pathogen after 72 hrs	Growth inhibition (%)
1	PTR-1	RS	3.4 ^g	62.2 ^d
2	PCF-3	RS	3.1 ^h	65.6 ^b
3	PTR-3	RS	2.8 ⁱ	68.9 ^a
4	PPF-2	RS	5.4 ^d	40 ^g
5	PPF-7	RS	5.1	43.3 ^f
6	PFC-1	RS	4.3 ^f	52.2 ^e
7	PTR-4	RS	4.1 ^f	54.4 ^d
8	PCF-7	RS	6.2 ^c	31.1 ^h
9	PFC-2	RS	6.9 ^b	23.3 ⁱ
10	PFC-7	RS	6.7 ^b	25.6 ^j
11	Control		9 ^a	

Data in the parentheses are arc sin means, data bearing same letter are not significantly different on the basis of DMRT at 5% level of significance

(0.10758 ug/ml). PTR-3 isolate showed highest P-solubilizing activity (17.3) followed by PTR-1 (14.2) (Table 5). Rassouli et al. (2005) reported that 201 indigenous *Pseudomonas* spp. isolated from Iranian soils were siderophore producers in CAS-agar medium. Kumar et al. (2000) reported that the *P. fluorescens* produced siderophores and antifungal metabolites which are involved in the control of phytopathogenic fungi. Gupta et al. (2002) reported *Pseudomonas* sp. acted as a potent phosphate solublizer. Correlation coefficients obtain between different biochemical parameters of rhizobacterial isolates and inhibitory activities against fungal pathogen (*R. solani*) were presented in Table 6 and Fig 1. The antagonistic activity of rhizobacterial isolates were found to be positively correlated with biochemical parameter of protease and chitinase activity. Among biochemical variables the correlation coefficient of chitinase and HCN production with antagonistic

potentiality of rhizobacteria was found to be significant at 1% level of significance. A linear multiple regression model was developed using different biochemical parameters through step wise technique can be used to predict the antagonistic activity of rhizobacterial isolates based on biochemical parameters. Using step wise regression technique a single biochemical variable (chitinase) was selected, which found to account for 94.0 % of variability of antagonistic potentiality and can be used as predictor of biocontrol potentiality of rhizobacteria on the basis of measured biochemical variables against fungal pathogens. Thus from the above finding it may be concluded that chitinase and protease production are the two most important biochemical variables significantly contributed towards the biocontrol potentialities of rhizobacteria against fungal pathogen (Equation 1). Biological control with fluorescent pseudomonad offers an effective

Table 3: Characterization of different isolates of rhizobacteria based on biochemical reaction

Isolates	Gram Reaction (3%KOH)	Growth at 4°C	Growth at 42°C	Arginine	Oxidase	catalase	Gelatin	trehlose	Levan	De-nitrification	Starch	citrate hydrolysis	Voges proskauer	Identified as
PTR-1	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PCF-3	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PTR-3	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PPF-2	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PFC-1	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PPF-7	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PFC-7	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PTR-4	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PBF-7	-	-	+	+	+	+	+	-	+	-	-	+	-	PA
PFC-2	-	-	+	+	+	+	+	-	+	-	-	+	-	PA

+: positive, - negative; PA: *Pseudomonas aurogenosa*

Table 4: Extracellular enzymes produced by rhizobacterial isolates

Isolates	Cellulase	Amalyse	Pectinase	Protease Halo zone (in cm)	Chitinase (p mol/s)
PTR-1	-	-	-	2.4 ^e	14.8 ^e
PCF-3	-	-	-	2.5 ^d	15.6 ^d
PTR-3	-	-	-	2.9 ^b	17.2 ^b
PPF-2	-	-	-	2.7 ^c	11.7 ^c
PPF-7	-	-	-	3 ^a	12.1 ^a
PFC-1	-	-	-	1.0 ⁱ	12.3 ⁱ
PTR-4	-	-	-	1.8 ^f	12.8 ^f
PCF-7	-	-	-	1.2 ^h	9.5 ^h
PFC-2	-	-	-	1.3 ^f	7.5 ^g
PFC-7	-	-	-	1.3 ^g	8.6 ^g

Data bearing same letter are not significantly different on the basis of DMRT at 5% level of significance

Table 5: Antimicrobial metabolites of different rhizobacterial isolates.

Isolates	IAA production (mg/ml)	HCN production (Absorbance at 625 nm)	Salicylic acid production (mg/ml)	Siderophore production (μ mole benzoic acid /ml)	Phosphate solubilization (mg/50 ml)
PTR-1	0.01039 ^d	0.0834 ^a	0.54 ^a	3.9 ^a	14.2 ^a
PCF-3	0.01023 ^d	0.0776 ^b	0.39 ^c	2.8 ^d	12.1 ^c
PTR-3	0.02677 ^e	0.0833 ^a	0.35 ^d	2.5 ^e	17.3 ^d
PPF-2	0.02786 ^e	0.0717 ^c	0.29 ^e	2.09 ^g	12.9 ^{ef}
PPF-7	0.06529 ^b	0.081 ^a	0.41 ^b	3.0 ^b	12.9 ^b
PFC-1	0.02688 ^e	0.0729 ^c	0.21 ^f	1.5 ^h	12.1 ^g
PTR-4	0.10758 ^a	0.0657 ^d	0.34 ^d	2.46 ^e	11.2 ^d
PCF-7	0.06351 ^b	0.0637 ^d	0.28 ^e	2.05 ^g	9.8 ^f
PFC-2	0.00207 ^e	0.034 ^e	0.22 ^f	2.19 ^f	12.3 ^e
PFC-7	0.00126 ^e	0.0324 ^e	0.19 ^g	2.42 ^e	10.3 ^d

Data bearing same letter are not significantly different on the basis of DMRT at 5% level of significance

Table 6: Correlation coefficient between antifungal antagonistic potentiality of rhizobacteria and their secondary metabolites production

Inhibition zone (Y)	Siderophore production (X1)	SA (X2)	HCN(X3)	IAA(X4)	P- Sol(X5)	Protease(X6)	Chitinase(X7)
Y	0.399	0.657*	0.823**	0.091	0.640*	0.557	0.980**




Figure 1: Relationship between chitinase activity and antibacterial antagonistic potentiality of rhizobacteria

method of managing plant pathogens (Ramamoorthy *et al.*, 2001). These bacteria inhibit the fungal pathogens by producing antibiotics, lytic enzymes and by inducing resistance systemically in the plant by activating defensive genes such as chitinase, α -1,3-glucanase, peroxidase and phenylalanine ammonia lyase (Frindlender *et al.*, 1991; Punja and Zhang, 1993; Ramamoorthy *et al.*, 2001; Viswanathan and Samiyappan, 2001). Chitinases are well known to lyse the fungal cell wall (Chet *et al.*, 1987). Addition of chitin or cell wall material to the fungus culture medium has been used in many studies to prove the chitinolytic activity (Ordentlich *et al.*, 1988; Frandberg and Schuurer, 1998; Viswanathan and Samiyappan, 2001).

Equation 1: Stepwise regression equation (secondary metabolites of rhizobacteria and their antifungal antagonistic potentiality)
 $Y = -17.367 + 5.244 (\text{Chitinase}); R^2 = 0.960; R^2 (\text{adj}) = 0.955;$
 $Y = -17.563 + 5.943 (\text{Chitinase}) - 4.147 (\text{protease}); R^2 = 0.980; R^2 (\text{adj}) = 0.974$

In conclusion, in this studies production of secondary metabolites such as chitinase, IAA, HCN, Salicylic Acid and siderophore by fluorescent *Pseudomonads* are important feature in plant disease suppression of root rot and enhancement of plant growth (Gade, 2013).

REFERENCES

- Adesemoye, A. O., Obini, M. and Ugoji, E. O. 2008. Comparison of plant growth promotion with *Pseudomonas aeruginosa* and *Bacillus subtilis* in three vegetables. *Braz. J. Mic.* **39**: 423-426.
- Alstrom, S. and Burns, R. G. 1989. Cyanide production by rhizobacteria as a possible mechanism of plant growth inhibition. *Biol. Fertil. Soils.* **7**: 232-238.
- Adhikari, A., Dutta, S., Nandi, S., Bhattacharya, I., De Roy, M., Sarkar, G. and Mandal, T. 2013. Antagonistic potentiality of native rhizobacterial isolates against root rot disease of okra, incited by *Rhizoctonia solani*. *African J. Agricultural Research.* **8(4)**: 405-412.
- Bossis, E. 1995. Les *Pseudomonas fluorescens* de la rhizosphère : étude taxonomique et effets sur la croissance de la tomate et du maïs, de la germination a la levee, These de doctorat, Universite de Nantes, France. p.143.
- Bossis, E., Lemanceau, P., Latour, X. and Gardan, L. 2000. The taxonomy of *Pseudomonas fluorescens* and *Pseudomonas putida*: current status and need for revision. *Agronomie.* **20**:51-63.
- Chet, L. 1987. *Trochoderma*-application, mode of action, and

potential as a biocontrol agent of soil borne plant pathogenic fungi. In: Innovative Approaches to Plant Disease Control. (Ed.) Chet, I. John Wiley and Sons, New York, pp. 137-160.

Frandberg, E. and Schuurer, I. 1998. Anti-fungal Activity of Chitinolytic Bacteria Isolated from Air Tight Stored Cereal Grain. *Can. J. Microbiol.*, **44**: 121-127.

Frindlender, M., Inbar, I. and Chet, L. 1991. Biological control of soil-borne plant pathogens by a α -1,3-glucanase producing *Pseudomonas cepacia*. *Soil Biol. Biochem.* **25**: 1211-1221.

Gade, R. M. 2013. Biological and chemical management of phytophthora root rot /collar rot in citrus nursery. *The Bioscan.* **7(4)**: 631-635, 2012

Gordon, S. A. and Weber, R. P. 1951. Calorimetric estimation of indoleacetic acid. *Plant growth-promoting rhizobacteria. Biocontrol Sci. and Techno.* **11**: 557-574.

Gupta, A., Meyer, J. M. and Goel, R. 2002. Development of heavy metal resistant mutants of phosphate solubilizing *Pseudomonas* sp. NBRI 4014 and their characterizaiton. *Curr. Microbiol.* **45**: 323-327.

ISTA 1966. International Rules of Seed Testing; *Proc. Int. Seed. Test. Assoc.* **32**: 565-589.

Jeon, J. S., Lee, S. S., Kim, H. Y., Ahn, T. S. and Song, H. G. 2003. Plant growth promotion in soil by some inoculated microorganisms. *J. Microbiol.* **41**: 271-276.

Johnson, L. F. and Curl, E. A. 1977. *Methods for Research on the Ecology of Soil Borne Plant Pathogens.* Burgess Publishing Company, Minneapolis.

Krishnaveni, M. 1991. Biological control of sheath blight disease of rice caused by *Rhizoctonia solani* Kuhn with *Pseudomonas fluorescens*. Masters Dissertation, University of Madras.

Kumar, G., Sharma, R. C. and Rai, S. N. 2000. Biocontrol of banded leaf and sheath blight of maize by peat based *Pseudomonas fluorescens* formulation. *Indian Phytopathol.* **53**: 190-192.

Lorck, H. 1948. Production of hydrocyanic acid by bacteria. *Physiol. Pl.* **1**: 142-146.

Luo JinYan., Xie GuanLin., Li Bin., Luo Yuan Chan., Zhao LiHan., Wang Xiao., Liu, B. and Li, W. 2005. Gram-positive bacteria associated with rice in China and their antagonists against the pathogens of sheath blight and bakanae disease in rice. *Rice Science.* **12(3)**: 213-218.

Meyer, J. M., Ajelvandire, P. and Georges, C. 1992. Iron metabolism in *Pseudomonas*: Salicylic acid, a siderophore of *Pseudomonas fluorescens* CHAO. *Biofactors.* **4**: 23-27.

Ordentlich, A., Elad, Y. and Chet, L. 1988. The role of chitinase of *Serratia marcescens* in Biocontrol of *Sclerotium rolfsii*. *Phytopath.* **78**: 84-88.

Mina, D. Koche, Gade, R. M. and Deshmukh, A. G. 2013. Antifungal activity of secondary metabolites produced by *pseudomonas fluorescens*. *The Bioscan.* **8(2)**: 723-726.

Punja, Z. K. and Zhang, Y. Y. 1993. Plant Chitinase and their roles in resistance to fungal disease. *J. Nematol.* **25**: 526-540.

Ramamoorthy, V., Viswanathan, R., Raguchander, T., Prakasam, V. and Samiyappan, R. 2001. Induction of Systemic Resistance by Plant Growth Promoting Rhizobacteria in Crop Plants against Pests and Diseases. *Crop Protect.* **20**: 1-11.

Rassouli, M. H., Khavazi, K., Rahimian, H., Malakouti, M. J., Asadi-Rahmani, H. 2005. An evaluation of the potential of indigenous Fluorescent *Pseudomonads* of wheat rhizosphere for producing siderophore. *Iran J. Soil Water Sci.* **20**:133-143.

Reddy, K. R. N. Choudary, D. A. and Reddy, M. S. 2007. Antifungal metabolites of *Pseudomonas fluorescens* isolated from rhizosphere of

Rice crop. *J. Mycol Pl Pathol.* **37**: 280-284.

Reeves, M., Pine, L., Neilands, J. B. and Bullows, A. 1983. Absence of siderophore activity in *Legionella* spp. grown in iron deficient media. *J. Bacteriol.* **154**: 324-329.

Smibert, R. M. and Krieg, N. R. 1994. Phenotypic characterization. In Gerharel P, Murray R GE, Wood WA, Krieg NR (Eds) Methods for general and molecular bacteriology. *American Society of Microbiology*, Washington, DC. pp. 607-654.

Stolp, H. and Gadkari, D. 1981. Non pathogenic members of genus *Pseudomonas*. In: The Prokaryotes: A Handbook on Habitats, Isolation and Identification of Bacteria (Eds. Starr, M.P., Stolp, H., Truper, H.G., Balows, A. and Schlegel, H.G.), Springer-Verlag, Berlin. I: 719-741.

Tripathi, M. and Johri, B. N. 2002. *In vitro* antagonistic potential of

fluorescent and control of sheath blight of maize caused by *Rhizoctonia solani*. *Indian J. Microbiol.* **42**: 207-214.

Vasantha Devi, T., Malarvizhi, R., Sakthivel, N. and Gnanamanickam, S. S. 1989. Biological control of sheath blight of rice in India with antagonistic bacteria. *Plant Soil.* **119**: 325-330.

Vessey, I. C. and Pegg, G. F. 1973. Autolysis and chitinases production in cultures *Verticillium albo-atrum*. *Transaction of the British Mycol. Soc.* **60**: 133-143.

Vincent, J. M. 1947. Distortion of fungal hyphae in the presence of certain inhibitors. *Nature.* **150**: 850.

Viswanathan and Samiyappan 2001. Antifungal Activity of Chitinase Produced by Fluorescent *Pseudomonads* against *Colletotrichum falcatum* Went Causing Redroot Disease in Sugarcane. *Microbiol Res.* **172**: 4017-4022.

NATIONAL ENVIRONMENTALISTS ASSOCIATION AND ITS OFFICIAL ORGAN



The Bioscan

An International Quarterly Journal of Life Science

Started in 1988, the National Environmentalists Association has been reorganized in 2006 and now is an association functioning with full vigour and new impetus to meet its objectives with the co-operation of like minded environment conscious academicians from different parts of the nation.

MEMBERSHIP OF THE ASSOCIATION

Any graduate having interest in environmental conservation and protection of nature and natural resources can be the member of the association.

To be the member of the association the application form given below should be duly filled up and sent to the Secretary of the association along with a demand draft of Rs. 750/- (After the 25% concession) for annual membership and Rs. 7500/- (After the 25% concession) for life membership.

FELLOWSHIP OF THE ASSOCIATION

The Association is awarding FELLOWSHIP to deserving academicians / researchers /scientists who are LIFE MEMBERS of the Association after reviewing their bio-data by the Fellows and the Executive Members of the association. The Fellows are privileged to write **F.N.E.A.** after their names .The prestigious Fellowship also includes a citation in recognition of their contribution to society in general and the endeavour for the noble cause of environment in particular.

AWARDS OF THE ASSOCIATION

The Association in its Seminars and Conferences provides the following category of awards on annual basis.

1. **The young scientists award** : It is given to the researchers below the age of 35 years.
2. **The senior scientists award** : It is awarded to the academicians above the age of 35 years.

3. **The best paper award**: It is awarded to the contributor of the Journal **The Bioscan** during the year.
4. **The best paper presentation award** : It is awarded to the scholar whose presentation is the best other than the young scientist category.
5. **The best oration award** : It is awarded to the scholar who delivered invited speech.
6. **The recognition award** : It is awarded to those senior scholars who have contributed to the subject through their continued research .
7. **The environmental awareness award** : It is awarded to those who, apart from their research contribution, have done commendable extension work for environmental betterment.

The number of recipients of award in each category will vary depending upon the recommendation of the panel of judges and the executive committee. The association has the provision to institute awards in the name of persons for whom a with desired sum is donated in consultation with the executive body.

PUBLICATION OF THE ASSOCIATION

In order to provide a platform to a vast group of researchers to express their views and finding of research as well as to promote the attitude of quality research among the scholars of younger generation the association publishes an international quarterly journal – **THE BIOSCAN (ISSN:0973-7049)**. For the benefit of the potential contributors **instructions to authors** is given separately in this journal. However, the details regarding the journal and also the association can be seen on our website www.thebioscan.in.

Cont. P. 270